



Representations of Homotopy group through infinite symmetric product functor

Pravanjan Kumar Rana[✉]

Assistant Professor and Head, Dept. of Mathematics, Berhampore Girls' College; Berhampore, Murshidabad, West Bengal, India

[✉]**Corresponding author:** Assistant Professor and Head, Dept. of Mathematics, Berhampore Girls' College; Berhampore, Murshidabad, West Bengal, India, Mail: - rana_pravanjan@yahoo.co.in

Publication History

Received: 04 August 2013

Accepted: 16 September 2013

Published: 1 October 2013

Citation

Pravanjan Kumar Rana. Representations of Homotopy group through infinite symmetric product functor. *Discovery*, 2013, 6(16), 7-9

Publication License



© The Author(s) 2013. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).

General Note

Article is recommended to print as color digital version in recycled paper.

ABSTRACT

In this paper we construct symmetric product covariant functor and we study the new representations of higher order Homotopy group through infinite symmetric product covariant functor(SP^∞) and higher order Homotopy functor(π_n , where $n \geq 0$).

Key words: Covariant functor, Natural transformation, Homotopy Equivalence, Homotopy Group.

1. INTRODUCTION

We recall the following definitions and statements

Definition 1.1

Let X be a topological space with a base point $x_0 \in X$. For $n \geq 0$, we define the n -fold symmetric product of X , denoted by $SP^n X$ by $SP^n X = X_0^n / S_n$ for $n \geq 1$, where X^n denotes the n -fold Cartesian product of X with itself and S_n denotes the symmetric group on n -objects regarded as acting on X^n by permuting the coordinates.

Hence for $n \geq 1$,

$$SP^n X = \{(x_1, x_2, x_3, \dots, x_n) : x_i \in X\}$$

Definition 1.2

Let X be a topological space with a base point $x_0 \in X$. Then there is a natural continuous map

$\mu: SP^n X \rightarrow SP^{n+1} X$ defined by

$$\mu(x_1, x_2, x_3, \dots, x_n) = (x_0, x_1, x_2, \dots, x_n),$$

$\Rightarrow \mu$ is a homeomorphism of $SP^n X$ onto the closed subset of $SP^{n+1} X$.

Thus we get the following sequence of spaces

$$x_0 = SP^0 X \subset SP^1 X \subset \dots \subset SP^n X \subset SP^{n+1} X \subset \dots$$

Thus we define $\lim_{n \rightarrow \infty} SP^n X = \bigcup_{n=1}^{\infty} SP^n X$, is called the infinite symmetric product of X and is denoted by $SP^\infty X$;

$$\text{i.e., } SP^\infty X = \bigcup_{n=1}^{\infty} SP^n X,$$

Definition 1.3

A category C consists of a class of objects X, Y, \dots , denoted by $Ob(C)$

a. for each ordered pair of objects X, Y a set of morphism with domain X and range Y denoted by $C(X, Y)$

- b. for each ordered triple of objects X, Y and Z and a pair of morphisms $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, their composite is denoted by $gf: X \rightarrow Z$, satisfying the two axioms:
- associativity
 - identity

Definition 1.4

Let \mathbf{C} and \mathbf{D} be two categories. A covariant functor T from \mathbf{C} to \mathbf{D} consists of

- an object function which assigns to every object X of \mathbf{C} and object $T(X)$ of \mathbf{D} ; and
 - a morphism function which assigns to every morphism $f: X \rightarrow Y$ in \mathbf{C} , a morphism $T(f): T(X) \rightarrow T(Y)$ in \mathbf{D} such that
- $T(1_X) = 1_{T(X)}$,
 - $T(gf) = T(g)T(f)$, for $g: Y \rightarrow Z$ in \mathbf{C}

Definition 1.5

Let \mathbf{C} and \mathbf{D} be categories. Suppose T_1 and T_2 are both covariant functors from \mathbf{C} to \mathbf{D} .

A natural transformation Φ from T_1 to T_2 is a function from the objects of \mathbf{C} to the morphisms of \mathbf{D} such that for every morphism $f: X \rightarrow Y$ in \mathbf{C} the following condition hold:

$$T_2(f)\Phi(X) = \Phi(Y)T_1(f)$$

Definition 1.6

Let X be a pointed topological spaces with base point x_0 and let the set $F_n(X, x_0)$ of all continuous maps α from I^n into X for which $\alpha(\partial I^n) = x_0$.

Define a relation \sim on $F_n(X, x_0)$ as follows:

for α and β in $F_n(X, x_0)$, $\alpha \sim \beta$, if there is a homotopy $H: I^n \times I \rightarrow X$

$$\text{such that } H(t_1, t_2, t_3, \dots, t_n, 0) = \alpha(t_1, t_2, t_3, \dots, t_n),$$

$$H(t_1, t_2, t_3, \dots, t_n, 1) = \beta(t_1, t_2, t_3, \dots, t_n), (t_1, t_2, t_3, \dots, t_n) \in I^n$$

$$H(t_1, t_2, t_3, \dots, t_n, s) = x_0, (t_1, t_2, t_3, \dots, t_n) \in \partial I^n, s \in I,$$

This relation \sim is an equivalence relation on $F_n(X, x_0)$, the equivalence class determined by α is denoted by $[\alpha]$, is called the homotopy class of α .

The set of homotopy classes of α is denoted by $\pi_n(X, x_0)$. The set of homotopy classes of α of $F_n(X, x_0)$ is a group. This group is called the n -th homotopy group of X at x_0 and is denoted by $\pi_n(X, x_0)$.

Definition 1.7


A quasifibration is a map $p: E \rightarrow B$ such that for all $b \in B$ and $e \in p^{-1}(b)$, $\rho_*: \pi_n(E, p^{-1}(b)) \rightarrow \pi_n(B, b)$ is an isomorphism, where the homotopy functors are based on e and b respectively

Definition 1.8

Let X be a path connected Hausdorff pointed space and let $A \subset X$. A neighbourhood $B \supset A$ is called deformable to A if there exists a homotopy $F: X \times I \rightarrow X$ such that

$$F(x, 0) = x, \text{ for all } x,$$

$$F(A \times I) \subseteq A, F(B \times I) \subseteq B \text{ and } F(B \times \{1\}) \subseteq A.$$



Definition 1.9

Let A and B be any two subsets of X . B is said to be deformable into A over X if the identity map $1_B: B \rightarrow B$ is homotopic in X to a map of B into A .

Lemma 1.10

If $f: X \rightarrow Y$ and $g: X \rightarrow Y$ are such that $f \simeq g$, then

- $SP^n f \simeq SP^n g$
- $SP^\infty f \simeq SP^\infty g$

Lemma 1.11

Let \mathcal{C} be any category and T_1, T_2 covariant functors from \mathcal{C} to \mathcal{C} . Then for any object C in \mathcal{C} , there is an equivalence $\theta: (T_1, T_2) \rightarrow T_2(C)$, where (T_1, T_2) is the class of natural transformations from T_1 to T_2 . This Lemma is called Yoneda's Lemma.

Lemma 1.12

If $f: X \rightarrow Y$ is a homotopy equivalence between two pointed topological spaces, then $SP^n f$ and $SP^\infty f$ are also homotopy equivalence.

2. STUDY OF SYMMETRIC PRODUCT FUNCTOR AND HOMOTOPY FUNCTOR

In this section we study the homotopy functor π_n associated with homotopy group $\pi_n(X, x_0)$ and a symmetric product covariant functor SP^n . To do this we use the following theorems:

Theorem 2.1 (Dold- Thom)

Let X be a Hausdorff pointed space and A be a closed path connected subspace with neighbourhood deformable to it. If $p: X \rightarrow X/A$ is the quotient map, then $SP^\infty p$ is a quasifibration with fibre $(SP^\infty p)^{-1}(x)$ homotopy equivalent to $SP^\infty A$ for all x in $SP^\infty X/A$.

Theorem 2.2

Let X be a Hausdorff pointed space such that Y is path connected with a map $f: X \rightarrow Y$.

Let $\rho: C_f \rightarrow \Sigma X$ be the quotient map from the mapping cone of f to the suspension of X which collapses $Y \times \{0\}$ to a point. Then $SP^\infty \rho: SP^\infty C_f \rightarrow SP^\infty \Sigma X$ is a quasifibration with fibre $SP^\infty X$.

Theorem 2.3

Let $p: E \rightarrow B$ be a quasifibration. Given $b \in B$ and $e \in p^{-1}(b)$ there exists a long exact sequence

$$\dots \rightarrow \pi_n(F) \rightarrow \pi_n(E) \rightarrow \pi_n(B) \rightarrow \pi_{n-1}(F) \rightarrow \dots$$

where $F = p^{-1}(b)$ is the fiber and $i: F \rightarrow E$ is the inclusion.

Proposition 2.4

Let X be a path connected Hausdorff pointed space. Then for all $n \geq 0$, $\pi_n(SP^\infty X) \cong \pi_{n+1}(SP^\infty \Sigma X)$. Also this isomorphism is natural.

Proof

Applying the **Theorem 2.2** to the identity map and noting that $C_{id} \cong CX$, the long exact sequence of the quasifibration $SP^\infty \rho: SP^\infty CX \rightarrow SP^\infty \Sigma X$ gives

$$\dots \rightarrow \pi_{n+1}(SP^\infty CX) \rightarrow \pi_{n+1}(SP^\infty \Sigma X) \rightarrow \pi_n(SP^\infty X) \rightarrow \pi_n(SP^\infty CX) \rightarrow \dots$$

Since CX is contractible, $SP^\infty CX$ is contractible and has trivial homotopy, so this segment of the sequence reduces to

$$\begin{aligned} 0 &\rightarrow \pi_{n+1}(SP^\infty \Sigma X) \rightarrow \pi_n(SP^\infty X) \rightarrow 0 \\ &\Rightarrow \pi_n(SP^\infty X) \cong \pi_{n+1}(SP^\infty \Sigma X). \text{ In particular we have} \\ 0 &\rightarrow \pi_1(SP^\infty \Sigma X) \rightarrow \pi_0(SP^\infty X) \cong 0 \\ &\Rightarrow \pi_1(SP^\infty \Sigma X) \cong 0 \end{aligned}$$

Next we show that this isomorphism is natural.

Given a map $f: X \rightarrow Y$, this is equivalent to that the right square of

Theorem 2.5

Let $f: (X, x_0) \rightarrow (Y, y_0)$ be a base point preserving continuous map then

$\pi_n(f): \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$ is a homomorphism.

Proof

The function $f_* = \pi_n(f): \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$ be defined by $f_*([\alpha]) = [f\alpha]$, $[\alpha] \in \pi_n(X, x_0)$.

Let $[\alpha], [\beta] \in \pi_n(X, x_0)$ then $f_*([\alpha] \circ [\beta]) = f_*([\alpha * \beta]) = [f(\alpha * \beta)] = [f\alpha] \circ [f\beta] = f_*([\alpha]) \circ f_*([\beta])$,

Thus $f_*([\alpha] \circ [\beta]) = f_*([\alpha]) \circ f_*([\beta])$

\Rightarrow The function f_* is a homomorphism.

All Homotopy groups and their homomorphisms from a category, this category will be denoted by '**Hop**' then we have the following theorem:

Theorem 2.6

For all $n \geq 0$, $\pi_n: \mathbf{Top} \rightarrow \mathbf{Hop}$ is a covariant functor.

Proof

Let (X, x_0) be the pointed topological spaces in **Top**, then $\pi_n(X, x_0)$ be the homotopy group in **Hop**.

Let $f: (X, x_0) \rightarrow (Y, y_0)$ and $g: (Y, y_0) \rightarrow (Z, z_0)$ are in **Top**, then

$f_* = \pi_n(f): \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$ be defined by

$f_*([\alpha]) = [f\alpha]$, $[\alpha] \in \pi_n(X, x_0)$.

Now $(g \circ f): (X, x_0) \rightarrow (Z, z_0)$ in '**Top**'.

Then using the **Theorem 2.5**, we have $\pi_n(g \circ f) = \pi_n(g) \circ \pi_n(f)$ and $\pi_n(l(X, x_0))$

$$= \pi_n(X, x_0)$$

Let all symmetric groups and their homomorphisms from a category, this category will be denoted by '**Sop**'.

Thus the theorem follows.

Theorem 2.7

$SP^n: \mathbf{Top} \rightarrow \mathbf{Sop}$ is a covariant functor, for all $n \in \mathbb{N}$

Proof

Let $f: X \rightarrow Y$ be base point preserving continuous map and

$SP^n(f): SP^n(X) \rightarrow SP^n(Y)$ be defined by

$SP^n f(x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)}) = (f(x_{\alpha(1)}), f(x_{\alpha(2)}), f(x_{\alpha(3)}), \dots, f(x_{\alpha(n)}))$

$SP^n l(x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)}) = (l(x_{\alpha(1)}), l(x_{\alpha(2)}), l(x_{\alpha(3)}), \dots, l(x_{\alpha(n)}))$

$= (x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)})$

i.e, $SP^n l(x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)}) = (x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)})$

$\Rightarrow SP^n l = Id$.

Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be two base point preserving continuous maps, then

$g \circ f: X \rightarrow Z$ is also base point preserving continuous map and $SP^n(g \circ f): SP^n X \rightarrow SP^n Z$ be defined by

$SP^n(g \circ f)(x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)})$

$= ((g \circ f)(x_{\alpha(1)}), (g \circ f)(x_{\alpha(2)}), (g \circ f)(x_{\alpha(3)}), \dots, (g \circ f)(x_{\alpha(n)}))$

$= (g(f(x_{\alpha(1)})), g(f(x_{\alpha(2)})), g(f(x_{\alpha(3)})), \dots, g(f(x_{\alpha(n)})))$

$= SP^n g(f(x_{\alpha(1)}), f(x_{\alpha(2)}), f(x_{\alpha(3)}), \dots, f(x_{\alpha(n)}))$

$= SP^n g \circ SP^n f(x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)})$

$$= (SP^n g \circ SP^n f)(x_{\alpha(1)}, x_{\alpha(2)}, x_{\alpha(3)}, \dots, x_{\alpha(n)})$$

Thus $SP^n(g \circ f) = SP^n g \circ SP^n f$

$\Rightarrow SP^n$ is a covariant functor

Corollary 2.8

SP^∞ is also a covariant functor

Proof

Using the **Definition 1.2** and **Theorem 2.7**, it follows.

3. NEW REPRESENTATION OF HIGHER ORDER HOMOTOPY GROUP ASSOCIATED WITH INFINITE SYMMETRIC PRODUCT FUNCTORS

Theorem 3.1

Let **Top** denotes the category of pointed topological space and base point preserving continuous maps and **Hop** denotes the category of Homotopy groups and their homomorphisms, then there is an equivalence $\theta: (\pi_n, SP^n) \rightarrow \pi_n(X, x_0)$, where (π_n, SP^n) denotes the set of all natural transformations from π_n to SP^n .

Proof

Using the **Definition 1.5** and **Lemma 1.11**, it follows.

Corollary 3.2

Let **Top** denotes the category of pointed topological space and base point preserving continuous maps and **Hop** denotes the category of Homotopy groups and their homomorphisms, then there is an equivalence $\theta: (\pi_n, SP^\infty) \rightarrow \pi_n(X, x_0)$, where (π_n, SP^∞) is the set of all natural transformations from π_n to SP^∞ .

Proof

Using **Theorem 3.1** and using $\lim_{n \rightarrow \infty} SP^n X = \bigcup_{n=1}^{\infty} SP^n X$, it follows.

Using the **Theorem 3.1**, we have the following:

Theorem 3.3

Let **Top** denotes the category of pointed topological space and base point preserving continuous maps and **Hop** denotes the category of Homotopy groups and their homomorphisms, then the cardinal number $|(\pi_n, SP^n)| = |\pi_n(X, x_0)|$, where π_n and SP^n are covariant functors;

where $|X|$ denote the underlying set of the topological space.

Theorem 3.4

Let **Top** denotes the category of pointed topological space and base point preserving continuous maps and **Hop** denotes the category of Homotopy groups and their homomorphisms, then the cardinal number $|(\pi_n, SP^\infty)| = |\pi_n(X, x_0)|$ where π_n and SP^∞ are covariant functors,

where $|X|$ denote the underlying set of the topological space.

REFERENCE

- Adhikary MR. Groups, Rings and Modules with Applications Universities Press, India, 1999
- Massey WS. Algebraic topology; an introduction, Springer Verlag, 1984
- Munkers JR. Topology a first course. Prentice Hall Inc, 1975
- Spanier H. Algebraic topology, Tata Mc-Graw-Hill Pub. Co. Ltd, 1966
- Adhikary A, Rana PK. A study of functors associated with Topological groups, Studia Univ. "Babes-Bolyai," Mathematica, Dec, 2001, Vol XLVI, 4
- Rana PK. A study of the group of covering transformation through functors, *BULLETIN MATHEMATIQUE*, 2009, 33(LIX), 21-24
- Rana PK. A study of functors associated with rings on continuous functions, *JIAM*, 2011, 33(1), 73-78
- Rana PK. A study of some homotopy type invariant functor. *International Journal of Science, Technology and Management*, 2012, 2, 67-71